

Table 5.15  
Cancer of the Cervix Uteri (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2011  
By Race/Ethnicity and Years Since Diagnosis

| Years Since Diagnosis               | 0 to <5 | 5 to <10 | 10 to <15 | 15 to <20 | 20 to <25 | 25 to <30 | 0 to <19 <sup>e</sup> | 0 to <36 <sup>e</sup> | >=36 <sup>g</sup> | Complete <sup>h</sup> |
|-------------------------------------|---------|----------|-----------|-----------|-----------|-----------|-----------------------|-----------------------|-------------------|-----------------------|
| <u>Race</u>                         |         |          |           |           |           |           |                       |                       |                   |                       |
| All Races <sup>b</sup>              | 39,578  | 33,185   | 35,754    | 32,230    | 28,157    | 20,702    | 147,326               | 211,309               | 38,323            | 249,632               |
| White <sup>b</sup>                  | 31,065  | 26,389   | 28,581    | 26,721    | 23,662    | 17,154    | 118,141               | 171,365               | 32,320            | 203,685               |
| Black <sup>b</sup>                  | 5,426   | 4,525    | 4,703     | 3,484     | 2,922     | 2,720     | 18,942                | 26,687                | +                 | +                     |
| Asian/Pacific Islander <sup>c</sup> | 2,147   | 1,764    | 1,663     | +         | +         | +         | 6,706                 | +                     | +                 | +                     |
| Hispanic <sup>d</sup>               | 7,909   | 7,382    | 6,634     | +         | +         | +         | 26,543                | +                     | +                 | +                     |

Estimated prevalence percent<sup>a</sup> on January 1, 2011, of the SEER<sup>c</sup> population diagnosed in the previous 19 years  
By Age at Prevalence and Race/Ethnicity

| Age at Prevalence                   | Age Specific (Crude) |     |       |         |         |         |         |         |         |         | Age-Adjusted <sup>f</sup> |
|-------------------------------------|----------------------|-----|-------|---------|---------|---------|---------|---------|---------|---------|---------------------------|
|                                     | All Ages             | 0-9 | 10-19 | 20-29   | 30-39   | 40-49   | 50-59   | 60-69   | 70-79   | 80+     | All Ages                  |
| <u>Race</u>                         |                      |     |       |         |         |         |         |         |         |         |                           |
| All Races <sup>c</sup>              | 0.0928%              | -   | -     | 0.0089% | 0.0762% | 0.1797% | 0.1983% | 0.1762% | 0.1544% | 0.0989% | 0.0879%                   |
| White <sup>c</sup>                  | 0.0966%              | -   | -     | 0.0106% | 0.0875% | 0.1982% | 0.2039% | 0.1663% | 0.1407% | 0.0802% | 0.0913%                   |
| Black <sup>c</sup>                  | 0.0789%              | -   | -     | 0.0055% | 0.0612% | 0.1402% | 0.1892% | 0.1900% | 0.1722% | 0.1539% | 0.0819%                   |
| Asian/Pacific Islander <sup>c</sup> | 0.0820%              | -   | -     | 0.0036% | 0.0325% | 0.1095% | 0.1693% | 0.2115% | 0.2172% | 0.1893% | 0.0755%                   |
| Hispanic <sup>d</sup>               | 0.1082%              | -   | -     | 0.0093% | 0.0806% | 0.2353% | 0.3394% | 0.3373% | 0.3350% | 0.2624% | 0.1410%                   |

<sup>a</sup> US 2011 cancer prevalence counts are based on 2011 cancer prevalence proportions from the SEER registries and 1/1/2011 US population estimates based on the average of 2010 and 2011 population estimates from the US Bureau of the Census.  
<sup>b c d</sup> Prevalence was calculated using the First Malignant Primary Only for a person. Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.  
<sup>e</sup> Maximum limited-duration prevalence: 36 years for 1975-2011 SEER 9 data; 19 years for 1992-2011 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).  
<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.  
<sup>g h i</sup> (g) Cases diagnosed more than 36 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <36 and >=36. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.  
- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.  
+ Not available.